



ESSENTIAL MEDICINES

**LIST @Doctors
Care**

@Doctors Care

1. ANAESTHETICS

1.1 General anaesthetics and oxygen

④ halothane	Inhalation.
ketamine	Injection: 50 mg (as hydrochloride)/ml in 10-ml vial.
nitrous oxide	Inhalation.
oxygen	Inhalation (medicinal gas).
④ thiopental	Powder for injection: 0.5 g; 1.0 g (sodium salt) in ampoule.

1.2 Local anaesthetics

④ bupivacaine	Injection: 0.25%; 0.5% (hydrochloride) in vial. Injection for spinal anaesthesia: 0.5% (hydrochloride) in 4-ml ampoule to be mixed with 7.5% glucose solution.
④ lidocaine	Injection: 1%; 2% (hydrochloride) in vial. Injection for spinal anaesthesia: 5% (hydrochloride) in 2-ml ampoule to be mixed with 7.5% glucose solution. Topical forms: 2-4% (hydrochloride).
lidocaine + epinephrine (adrenaline)	Dental cartridge: 2% (hydrochloride) + epinephrine 1:80 000. Injection: 1%; 2% (hydrochloride) + epinephrine 1:200 000 in vial.

Complementary List

ephedrine	Injection: 30 mg (hydrochloride)/ml in 1-ml ampoule. <i>(For use in spinal anaesthesia during delivery, to prevent hypotension).</i>
-----------	--

1.3 Preoperative medication and sedation for short-term procedures

atropine	Injection: 1 mg (sulfate) in 1-ml ampoule.
④ diazepam	Injection: 5 mg/ml in 2-ml ampoule. Tablet: 5 mg.
morphine	Injection: 10 mg (sulfate or hydrochloride) in 1-ml ampoule.
promethazine	Oral liquid: 5 mg (hydrochloride)/5 ml.

ANTI-INFLAMMATORY MEDICINES (NSAIDs), MEDICINES USED TO TREAT GOUT AND DISEASE MODIFYING AGENTS IN RHEUMATOID DISORDERS (DMARDs)

2.1 Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs)

acetylsalicylic acid	Suppository: 50-150 mg. Tablet: 100-500 mg.
ibuprofen	Tablet: 200 mg; 400 mg.
paracetamol*	Oral liquid: 125 mg/5 ml. Suppository: 100 mg. Tablet: 100-500 mg. * Not recommended for anti-inflammatory use due to lack of proven benefit to that effect.
diclofenac (as Sodium)	75 mg / 3 ml inj. (as sodium)

2.2 Opioid analgesics

codeine	Tablet: 30 mg (phosphate).
morphine	Injection: 10 mg (morphine hydrochloride or morphine sulfate) in 1-ml ampoule. Oral liquid: 10 mg (morphine hydrochloride or morphine sulfate)/5 ml. Tablet: 10 mg (morphine sulfate). Tablet (prolonged release): 10 mg; 30 mg; 60 mg (morphine sulfate).

Pethidine	50 mg/ml inj. (as hydrochloride)
-----------	----------------------------------

2.3 Medicines used to treat gout

allopurinol	Tablet: 100 mg.
-------------	-----------------

2.4 Disease modifying agents used in rheumatoid disorders (DMARDs)

chloroquine	Tablet: 100 mg; 150 mg (as phosphate or sulfate).
-------------	---

Complementary List

azathioprine	Tablet: 50 mg.
methotrexate	Tablet: 2.5 mg (as sodium salt).
penicillamine	Capsule or tablet: 250 mg.
sulfasalazine	Tablet: 500 mg.

@Doctors
Care

2. ANALGESICS, ANTIPYRETICS, NON-STEROIDAL ANTI-INFLAMMATORY MEDICINES (NSAIMs), MEDICINES USED TO TREAT GOUT AND DISEASE MODIFYING AGENTS IN RHEUMATOID DISORDERS (DMARDs)

2.1 Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs)

acetylsalicylic acid	Suppository: 50-150 mg. Tablet: 100-500 mg.
ibuprofen	Tablet: 200 mg; 400 mg.
paracetamol*	Oral liquid: 125 mg/5 ml. Suppository: 100 mg. Tablet: 100-500 mg. * Not recommended for anti-inflammatory use due to lack of proven benefit to that effect.
diclofenac (as Sodium)	75 mg / 3 ml inj. (as sodium)

2.2 Opioid analgesics

codeine	Tablet: 30 mg (phosphate).
morphine	Injection: 10 mg (morphine hydrochloride or morphine sulfate) in 1-ml ampoule. Oral liquid: 10 mg (morphine hydrochloride or morphine sulfate)/5 ml. Tablet: 10 mg (morphine sulfate). Tablet (prolonged release): 10 mg; 30 mg; 60 mg (morphine sulfate).
Pethidine	50 mg/ml inj. (as hydrochloride)

2.3 Medicines used to treat gout

allopurinol	Tablet: 100 mg.
-------------	-----------------

2.4 Disease modifying agents used in rheumatoid disorders (DMARDs)

chloroquine	Tablet: 100 mg; 150 mg (as phosphate or sulfate).
-------------	---

Complementary List

azathioprine	Tablet: 50 mg.
methotrexate	Tablet: 2.5 mg (as sodium salt).
penicillamine	Capsule or tablet: 250 mg.
sulfasalazine	Tablet: 500 mg.

3. ANTIALLERGICS AND MEDICINES USED IN ANAPHYLAXIS

©chlorphenamine	Injection: 10 mg (hydrogen maleate) in 1-ml ampoule. Tablet: 4 mg (hydrogen maleate).
dexamethasone	Injection: 4 mg dexamethasone phosphate (as disodium salt) in 1-ml ampoule.
epinephrine (adrenaline)	Injection: 1 mg (as hydrochloride or hydrogen tartrate) in 1-ml ampoule.
hydrocortisone	Powder for injection: 100 mg (as sodium succinate) in vial.
© prednisolone*	Tablet: 5 mg; 25 mg * There is no evidence for complete clinical similarity between prednisolone and dexamethasone at high doses.

4. ANTIDOTES AND OTHER SUBSTANCES USED IN POISONINGS

4.1 Non-specific

charcoal, activated	Powder.
---------------------	----------------

4.2 Specific

acetylcysteine	Injection: 200 mg/ml in 10-ml ampoule.
atropine	Injection: 1 mg (sulfate) in 1-ml ampoule.
calcium gluconate	Injection: 100 mg/ml in 10-ml ampoule.
deferoxamine	Powder for injection: 500 mg (mesilate) in vial.
dimercaprol	Injection in oil: 50 mg/ml in 2-ml ampoule.
DL-methionine	Tablet: 250 mg.
methylthioninium chloride (methylene blue)	Injection: 10 mg/ml in 10-ml ampoule.
naloxone	Injection: 400 micrograms (hydrochloride) in 1-ml ampoule.
penicillamine	Capsule or tablet: 250 mg.
potassium ferric hexacyano-ferrate(II) -2H ₂ O (Prussian blue)	Powder for oral administration.
sodium calcium edetate	Injection: 200 mg/ml in 5-ml ampoule.

sodium nitrite	Injection: 30 mg/ml in 10-ml ampoule.
sodium thiosulfate	Injection: 250 mg/ml in 50-ml ampoule.

5. ANTICONVULSANTS/ANTIEPILEPTICS

carbamazepine	<p>Oral liquid: 100 mg/5 ml.</p> <p>Tablet (chewable): 100 mg; 200 mg.</p> <p>Tablet (scored): 100 mg; 200 mg.</p>
① diazepam	Injection: 5 mg/ml in 2-ml ampoule (intravenous or rectal).
magnesium sulfate*	<p>Injection: 500 mg/ml in 2-ml ampoule; 500 mg/ml in 10-ml ampoule.</p> <p>* For use in eclampsia and severe pre-eclampsia and not for other convulsant disorders.</p>
phenobarbital	<p>Injection: 200 mg/ml (phenobarbital sodium).</p> <p>Oral liquid: 15 mg/5 ml (as phenobarbital or phenobarbital sodium).</p> <p>Tablet: 15-100 mg (phenobarbital).</p>
phenytoin	<p>Capsule: 25 mg; 50 mg; 100 mg (sodium salt).</p> <p>Injection: 50 mg/ml in 5-ml vial (sodium salt).</p> <p>Oral liquid: 25 - 30 mg/5 ml.*</p> <p>Tablet: 25 mg; 50 mg; 100 mg (sodium salt).</p> <p>Tablet (chewable): 50 mg.</p> <p>* The presence of both 25 mg/5 ml and 30 mg/5 ml strengths on the same market would cause confusion in prescribing and dispensing and should be avoided.</p>
valproic acid	<p>Oral liquid: 200 mg/5 ml.</p> <p>Tablet (crushable): 100 mg.</p> <p>Tablet (enteric-coated): 200 mg; 500 mg (sodium valproate).</p>

Complementary List

ethosuximide	<p>Capsule: 250 mg.</p> <p>Oral liquid: 250 mg/5 ml.</p>
--------------	--

6.1 Anthelmintics

6.1.1 Intestinal anthelmintics

albendazole	Tablet (chewable): 400 mg.
levamisole	Tablet: 50 mg; 150 mg (as hydrochloride).
© mebendazole	Tablet (chewable): 100 mg; 500 mg.
niclosamide*	Tablet (chewable): 500 mg.
pyrantel	Oral liquid: 50 mg (as embonate)/ml. Tablet (chewable): 250 mg (as embonate).
ivermectin	Tablet (scored): 3 mg; 6 mg.

**@Doctors
Care**

6.2 Antibacterials

6.2.1 Beta Lactam medicines

amoxicillin	Capsule or tablet: 250 mg; 500 mg (anhydrous). Powder for oral liquid: 125 mg (anhydrous)/5 ml.
amoxicillin + clavulanic acid	Tablet: 500 mg + 125 mg.
ampicillin	Powder for injection: 500 mg; 1 g (as sodium salt) in vial.
benzathine benzylpenicillin	Powder for injection: 1.44 g benzylpenicillin (=2.4 million IU) in 5-ml vial.
benzylpenicillin	Powder for injection: 600 mg (= 1 million IU); 3 g (= 5 million IU) (sodium or potassium salt) in vial.
cefazolin*	Powder for injection: 1 g (as sodium salt) in vial. * For surgical prophylaxis.
cefixime*	Capsule: 400 mg. * Only listed for single-dose treatment of uncomplicated ano-genital gonorrhoea.
© cloxacillin	Capsule: 500 mg; 1 g (as sodium salt). Powder for injection: 500 mg (as sodium salt) in vial. Powder for oral liquid: 125 mg (as sodium salt)/5 ml.

phenoxycephalothin	Powder for oral liquid: 250 mg (as potassium salt)/5 ml. Tablet: 250 mg (as potassium salt).
procaine benzylpenicillin	Powder for injection: 1 g (=1 million IU); 3 g (=3 million IU) in vial.

Complementary List

ceftazidime	Powder for injection: 250 mg (as pentahydrate) in vial,
④ ceftriaxone	Powder for injection: 250 mg, 1 g (as sodium salt) in vial.
imipenem* + cilastatin*	Powder for injection: 250 mg (as monohydrate) + 250 mg (as sodium salt); 500 mg (as monohydrate) + 500 mg (as sodium salt) in vial. * Only listed for the treatment of life-threatening hospital-based infection due to suspected or proven multidrug-resistant infection.

6.2.2 Other antibacterials

azithromycin*	Capsule: 250 mg or 500 mg. Oral liquid: 200 mg/5 ml. * Only listed for single-dose treatment of genital <i>Chlamydia trachomatis</i> and of trachoma.
chloramphenicol	Capsule: 250 mg. Oily suspension for injection: 0.5 g (as sodium succinate)/ml in 2-ml ampoule. Oral liquid: 150 mg (as palmitate)/5 ml. Powder for injection: 1 g (sodium succinate) in vial.
④ ciprofloxacin*	Tablet: 250 mg (as hydrochloride). * Final selection depends on indication for use.
doxycycline*	Capsule or tablet: 100 mg (hydrochloride). * Final selection depends on indication for use.
④ erythromycin	Capsule or tablet: 250 mg (as stearate or ethyl succinate). Powder for injection: 500 mg (as lactobionate) in vial. Powder for oral liquid: 125 mg/ 5 ml (as stearate or ethyl succinate).
④ gentamicin*	Injection: 10 mg; 40 mg (as sulfate)/ml in 2-ml vial. * Final selection depends on indication for use.

©metronidazole	Injection: 500 mg in 100-ml vial. Oral liquid: 200 mg (as benzoate)/5 ml. Suppository: 500 mg; 1 g. Tablet: 200-500 mg.
nitrofurantoin	Tablet: 100 mg.
spectinomycin	Powder for injection: 2 g (as hydrochloride) in vial.
sulfamethoxazole + trimethoprim	Injection: 80 mg + 16 mg/ml in 5-ml and 10-ml ampoules. Oral liquid: 200 mg + 40 mg/5 ml. Tablet: 100 mg + 20 mg; 400 mg + 80 mg.
trimethoprim	Tablet: 100 mg; 200 mg.
nalidixic acid	500 mg tablet, 300 mg / 5ml syrup. 5000,000 IU tab, 100,000 IU/ml drops, 100,000 IU pessaries

Complementary List

clindamycin	Capsule: 150 mg. Injection: 150 mg (as phosphate)/ml.
sulfadiazine	Injection: 250 mg (sodium salt) in 4-ml ampoule. Tablet: 500 mg.

vancomycin	Powder for injection: 250 mg (as hydrochloride) in vial.
------------	---

6.2.3 Antileprosy medicines

Medicines used in the treatment of leprosy should never be used except in combination. Combination therapy is essential to prevent the emergence of drug resistance.

clofazimine	Capsule: 50 mg; 100 mg.
dapsone	Tablet: 25 mg; 50 mg; 100 mg.
rifampicin	Capsule or tablet: 150 mg; 300 mg.

6.2.4 Antituberculosis medicines

ethambutol	Tablet: 100-400 mg (hydrochloride).
------------	--

isoniazid	Tablet: 100-300 mg. Tablet (scored): 50 mg.
isoniazid + ethambutol	Tablet: 150 mg + 400 mg.
pyrazinamide	Tablet: 400 mg. Tablet (dispersible): 150 mg. Tablet (scored): 150 mg.
rifampicin	Capsule or tablet: 150 mg; 300 mg.
rifampicin + isoniazid	Tablet: 60 mg + 30 mg; 150 mg + 75 mg; 300 mg + 150 mg. 60 mg + 60 mg (For intermittent use three times weekly). 150 mg + 150 mg (For intermittent use three times weekly).
rifampicin + isoniazid + ethambutol	Tablet: 150 mg + 75 mg + 275 mg.
rifampicin + isoniazid + pyrazinamide	Tablet: 60 mg + 30 mg + 150 mg; 150 mg + 75 mg + 400 mg. 150 mg + 150 mg + 500 mg (For intermittent use three times weekly).
rifampicin + isoniazid + pyrazinamide + ethambutol	Tablet: 150 mg + 75 mg + 400 mg + 275 mg.
streptomycin	Powder for injection: 1 g (as sulfate) in vial.

Complementary List

amikacin	Powder for injection: 1000 mg in vial.
p-aminosalicylic acid	Granules: 4 g in sachet. Tablet: 500 mg.
capreomycin	Powder for injection: 1000 mg in vial.
cycloserine	Capsule or tablet: 250 mg.
ethionamide	Tablet: 125 mg; 250 mg.

<i>kanamycin</i>	<i>Powder for injection: 1000 mg in vial.</i>
<i>ofloxacin*</i>	<i>Tablet: 200 mg; 400 mg.</i> * <i>Levofloxacin may be an alternative based on availability and programme considerations.</i>
6.3 Antifungal medicines	
<i>clotrimazole</i>	<i>Vaginal cream: 1%; 10%.</i> <i>Vaginal tablet: 100 mg; 500 mg.</i>
<i>© fluconazole</i>	<i>Capsule: 50 mg.</i> <i>Injection: 2 mg/ml in vial.</i> <i>Oral liquid: 50 mg/5 ml.</i>
<i>griseofulvin</i>	<i>Capsule or tablet: 125 mg; 250 mg.</i>
<i>nystatin</i>	<i>Lozenge: 100 000 IU.</i> <i>Pessary: 100 000 IU.</i> <i>Tablet: 100 000 IU; 500 000 IU.</i>
Complementary List <i>amphotericin B</i> <i>Powder for injection: 50 mg in vial.</i> <i>flucytosine</i> <i>Capsule: 250 mg.</i> <i>Infusion: 2.5 g in 250 ml.</i> <i>potassium iodide</i> <i>Saturated solution.</i>	
6.4 Antiviral medicines	
6.4.1 Antiherpes medicines	
<i>© aciclovir</i>	<i>Powder for injection: 250 mg (as sodium salt) in vial.</i> <i>Tablet: 200 mg.</i>
6.4.2 Antiretrovirals	
6.4.2.1 Nucleoside/Nucleotide reverse transcriptase inhibitors	
<i>abacavir (ABC)</i>	<i>Oral liquid: 100 mg (as sulfate)/5 ml.</i> <i>Tablet: 300 mg (as sulfate).</i>

**@Doctors
Care**

didanosine (ddI)	Buffered powder for oral liquid: 100 mg; 167 mg; 250 mg packets. Capsule (unbuffered enteric-coated): 125 mg; 200 mg; 250 mg; 400 mg. Tablet (buffered chewable, dispersible): 25 mg; 50 mg; 100 mg; 150 mg; 200 mg.
emtricitabine (FTC)*	Capsule: 200 mg. Oral liquid: 10 mg/ml. * FTC is an acceptable alternative to 3TC, based on knowledge of the pharmacology, the resistance patterns and clinical trials of antiretrovirals.
lamivudine (3TC)	Oral liquid: 50 mg/5 ml. Tablet: 150 mg.
stavudine (d4T)	Capsule: 15 mg; 20 mg; 30 mg; 40 mg.* * The Committee expects this dosage form to be reviewed for possible deletion at the next meeting. Powder for oral liquid: 5 mg/5 ml.
tenofovir disoproxil fumarate (TDF)	Tablet: 300 mg (tenofovir disoproxil fumarate - equivalent to 245 mg tenofovir disoproxil).
zidovudine (ZDV or AZT)	Capsule: 100 mg; 250 mg. Oral liquid: 50 mg/5 ml. Solution for IV infusion injection: 10 mg/ml in 20-ml vial. Tablet: 300 mg.

6.4.2.2 Non-nucleoside reverse transcriptase inhibitors

efavirenz (EFV or EFZ)	Capsule: 50 mg; 100 mg; 200 mg. Oral liquid: 150 mg/5 ml. Tablet: 600 mg.
nevirapine (NVP)	Oral liquid: 50 mg/5 ml. Tablet: 200 mg.

6.4.2.3 Protease inhibitors

Ritonavir is recommended for use in combination as a pharmacological booster, and not as an antiretroviral in its own right.

indinavir (IDV)	Capsule: 200 mg; 333 mg; 400 mg (as sulfate).
lopinavir + ritonavir (LPV/r)	Capsule: 133.3 mg + 33.3 mg. Oral liquid: 400 mg + 100 mg/5 ml.
nelfinavir (NFV)	Oral powder: 50 mg/g. Tablet: 250 mg (as mesilate).
ritonavir	Oral liquid: 400 mg/5 ml. Oral solid dosage form: 100 mg.
saquinavir (SQV)	Capsule: 200 mg.

FIXED-DOSE COMBINATIONS

efavirenz + emtricitabine* + tenofovir	Tablet: 600 mg + 200 mg + 300 mg. * FTC is an acceptable alternative to 3TC, based on knowledge of the pharmacology, the resistance patterns and clinical trials of antiretrovirals.
emtricitabine* + tenofovir	Tablet: 200 mg + 300 mg. * FTC is an acceptable alternative to 3TC, based on knowledge of the pharmacology, the resistance patterns and clinical trials of antiretrovirals.
stavudine + lamivudine + nevirapine	Tablet: 30 mg + 150 mg + 200 mg.
zidovudine + lamivudine	Tablet: 300 mg + 150 mg.

6.4.3 Other antivirals

ribavirin	Injection for intravenous administration: 1000 mg and 800 mg in 10-ml phosphate buffer solution. Oral solid dosage forms: 200 mg; 400 mg; 600 mg.
-----------	--

6.5 Antiprotozoal medicines

diloxanide	Tablet: 500 mg (furoate).
------------	---------------------------

© metronidazole	Injection: 500 mg in 100-ml vial. Oral liquid: 200 mg (as benzoate)/5 ml. Tablet: 200-500 mg.
-----------------	---

6.5.2 Antileishmaniasis medicines

sodium stibogluconate (S)	100 mg inj.
------------------------------	-------------

Complementary List

amphotericin B	Powder for injection: 50 mg in vial.
----------------	--------------------------------------

6.5.3 Antimalarial medicines

6.5.3.1 For curative treatment

Medicines for the treatment of *P. falciparum* malaria cases should be used in combination.

amodiaquine*	Tablet: 153 mg or 200 mg (as hydrochloride). * To be used (a) in combination with artesunate 50 mg OR (b) may be used alone for the treatment of <i>P.vivax</i> , <i>P.ovale</i> and <i>P.malariae</i> infections.
artemether	Oily injection: 80 mg/ml in 1-ml ampoule. For use in the management of severe malaria.
artemether + lumefantrine*	Tablet: 20 mg + 120 mg. * Not recommended in the first trimester of pregnancy or in children below 5 kg.
artesunate*	Injection: ampoules, containing 60 mg anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution. For use in the management of severe malaria. Tablet: 50 mg. * To be used in combination with either amodiaquine, mefloquine or sulfadoxine + pyrimethamine.
chloroquine*	Oral liquid: 50 mg (as phosphate or sulfate)/5 ml. Tablet: 100 mg; 150 mg (as phosphate or sulfate). * For use only for the treatment of <i>P.vivax</i> infection.

**@Doctors
Care**

doxycycline*	Capsule: 100 mg (as hydrochloride). Tablet (dispersible): 100 mg (as monohydrate). * For use only in combination with quinine.
mefloquine*	Tablet: 250 mg (as hydrochloride). * To be used in combination with artesunate 50 mg.
primaquine*	Tablet: 7.5 mg; 15 mg (as diphosphate) * Only for use to achieve radical cure of <i>P.vivax</i> and <i>P.ovale</i> infections, given for 14 days.
quinine*	Injection: 300 mg quinine hydrochloride/ml in 2-ml ampoule. Tablet: 300 mg (quinine sulfate) or 300 mg (quinine bisulfate). * For use only in the management of severe malaria, and should be used in combination with doxycycline.
sulfadoxine + Pyrimethamine*	Tablet: 500 mg + 25 mg. * Only in combination with artesunate 50 mg.

6.5.3.2 For prophylaxis

chloroquine*	Oral liquid: 50 mg (as phosphate or sulfate)/5 ml. Tablet: 150 mg (as phosphate or sulfate). * For use only in central American regions, for use for <i>P.vivax</i> .
doxycycline	Capsule or tablet: 100 mg (hydrochloride).
mefloquine	Tablet: 250 mg (as hydrochloride).
proguanil*	Tablet: 100 mg (hydrochloride). * For use only in combination with chloroquine.

6.5.4 Antipneumocystosis and antitoxoplasmosis medicines

pyrimethamine	Tablet: 25 mg.
sulfamethoxazole + trimethoprim	Injection: 80 mg + 16 mg/ml in 5-ml ampoule; 80 mg + 16 mg/ml in 10-ml ampoule.

7. ANTIMIGRAINE MEDICINES

7.1 For treatment of acute attack

acetylsalicylic acid	Tablet: 300-500 mg.
----------------------	----------------------------

paracetamol	Tablet: 300-500 mg.
7.2 For prophylaxis	
© propranolol	Tablet: 20 mg; 40 mg (hydrochloride).
8. ANTINEOPLASTIC, IMMUNOSUPPRESSIVES	
8.1 Immunosuppressive medicines	
<i>Complementary List</i>	
azathioprine	<i>Powder for injection:</i> 100 mg (as sodium salt) in vial. <i>Tablet:</i> 50 mg.
ciclosporin	<i>Capsule:</i> 25 mg. <i>Concentrate for injection:</i> 50 mg/ml in 1-ml ampoule for organ transplantation.
8.2 Cytotoxic medicines	
<i>Complementary List</i>	
asparaginase	<i>Powder for injection:</i> 10 000 IU in vial.
bleomycin	<i>Powder for injection:</i> 15 mg (as sulfate) in vial.
calcium folinate	<i>Injection:</i> 3 mg/ml in 10-ml ampoule. <i>Tablet:</i> 15 mg.
chlorambucil	<i>Tablet:</i> 2 mg.
cisplatin	<i>Powder for injection:</i> 10 mg; 50 mg in vial.
cyclophosphamide	<i>Powder for injection:</i> 500 mg in vial. <i>Tablet:</i> 25 mg.
cytarabine	<i>Powder for injection:</i> 100 mg in vial.
dacarbazine	<i>Powder for injection:</i> 100 mg in vial.
dactinomycin	<i>Powder for injection:</i> 500 micrograms in vial.
daunorubicin	<i>Powder for injection:</i> 50 mg (as hydrochloride) in vial.
doxorubicin	<i>Powder for injection:</i> 10 mg; 50 mg (hydrochloride) in vial.
etoposide	<i>Capsule:</i> 100 mg. <i>Injection:</i> 20 mg/ml in 5-ml ampoule.
fluorouracil	<i>Injection:</i> 50 mg/ml in 5-ml ampoule.

<i>mercaptopurine</i>	<i>Tablet: 50 mg.</i>
<i>methotrexate</i>	<i>Powder for injection: 50 mg (as sodium salt) in vial. Tablet: 2.5 mg (as sodium salt).</i>
<i>procarbazine</i>	<i>Capsule: 50 mg (as hydrochloride).</i>
<i>vinblastine</i>	<i>Powder for injection: 10 mg (sulfate) in vial.</i>
<i>vincristine</i>	<i>Powder for injection: 1 mg; 5 mg (sulfate) in vial.</i>

@Doctors Care

8.3 Hormones and antihormones

Complementary List

<i>dexamethasone</i>	<i>Injection: 4 mg dexamethasone phosphate (as disodium salt) in 1-ml ampoule.</i>
<i>hydrocortisone</i>	<i>Powder for injection: 100 mg (as sodium succinate) in vial.</i>
<i>④ prednisolone*</i>	<i>Tablet: 5 mg; 25 mg.</i> <i>* There is no evidence for complete clinical similarity between prednisolone and dexamethasone at high doses.</i>
<i>tamoxifen</i>	<i>Tablet: 10 mg; 20 mg (as citrate).</i>

9. ANTIPARKINSONISM MEDICINES

<i>procyclidine (S), (T)</i>	<i>5 mg tab.</i>
<i>levodopa + ④</i>	<i>Tablet: 100 mg + 10 mg; 250 mg + 25 mg.</i>

10. MEDICINES AFFECTING THE BLOOD

10.1 Antianaemia medicines

<i>ferrous salt</i>	<i>Oral liquid: equivalent to 25 mg iron (as sulfate)/ml. Tablet: equivalent to 60 mg iron.</i>
<i>ferrous salt + folic acid</i>	<i>Tablet equivalent to 60 mg iron + 400 micrograms folic acid (Nutritional supplement for use during pregnancy).</i>
<i>folic acid</i>	<i>Tablet: 1 mg; 5 mg.</i>

<i>hydroxocobalamin</i>	<i>Injection: 1 mg in 1-ml ampoule.</i>
-------------------------	---

10.2 Medicines affecting coagulation

heparin sodium	Injection: 1000 IU/ml; 5000 IU/ml; 20,000 IU/ml in 1-ml ampoule.
phytomenadione	Injection: 10 mg/ml in 5-ml ampoule. Tablet: 10 mg.
protamine sulfate	Injection: 10 mg/ml in 5-ml ampoule.
© warfarin	Tablet: 1 mg; 2 mg; 5 mg (sodium salt).

11. BLOOD PRODUCTS AND PLASMA SUBSTITUTES

11.1 Plasma substitutes

© dextran 70*	Injectable solution: 6%. * Polygeline, injectable solution, 3.5% is considered as equivalent.
---------------	--

11.2 Plasma fractions for specific use

All plasma fractions should comply with the WHO Requirements for the Collection, Processing and Quality Control of Blood, Blood Components and Plasma Derivatives.

Complementary List

human normal immunoglobulin	<i>Intramuscular administration:</i> 16% protein solution. <i>Intravenous administration:</i> 5%, 10% protein solution.
© factor VIII concentrate	<i>Dried.</i>
© factor IX complex (coagulation factors, II, VII, IX, X) concentrate	<i>Dried.</i>

12. CARDIOVASCULAR MEDICINES

12.1 Antianginal medicines

© atenolol	Tablet: 50 mg; 100 mg.
glyceryl trinitrate	Tablet (sublingual): 500 micrograms.
© isosorbide dinitrate	Tablet (sublingual): 5 mg.
verapamil	Tablet: 40 mg; 80 mg (hydrochloride).

12.2 Antiarrhythmic medicines

This subsection will be reviewed at the next meeting of the Expert Committee.

① atenolol	Tablet: 50 mg; 100 mg.
digoxin	Injection: 250 micrograms/ml in 2-ml ampoule. Oral liquid: 50 micrograms/ml. Tablet: 62.5 micrograms; 250 micrograms.
epinephrine (adrenaline)	Injection: 100 micrograms/ml (as acid tartrate or hydrochloride) in 10-ml ampoule.
lidocaine	Injection: 20 mg (hydrochloride)/ml in 5-ml ampoule.

verapamil	Injection: 2.5 mg (hydrochloride)/ml in 2-ml ampoule. Tablet: 40 mg; 80 mg (hydrochloride).
-----------	--

Complementary List

① procainamide	Injection: 100 mg (hydrochloride)/ml in 10-ml ampoule.
① quinidine	Tablet: 200 mg (sulfate).

12.3 Antihypertensive medicines

**@Doctors
Care**

① amlodipine	Tablet: 5 mg.
① atenolol	Tablet: 50 mg; 100 mg.
① enalapril	Tablet: 2.5 mg.
hydralazine*	Powder for injection: 20 mg (hydrochloride) in ampoule. Tablet: 25 mg, 50 mg (hydrochloride). * Hydralazine is listed for use in the acute management of severe pregnancy-induced hypertension only. Its use in the treatment of essential hypertension is not recommended in view of the availability of more evidence of efficacy and safety of other medicines.
① hydrochlorothiazide	Tablet (scored): 25 mg.

methyldopa*	Tablet: 250 mg. * Methyldopa is listed for use in the management of pregnancy-induced hypertension only. Its use in the treatment of essential hypertension is not recommended in view of the availability of more evidence of efficacy and safety of other medicines.
-------------	--

Complementary List

sodium nitroprusside	<i>Powder for infusion:</i> 50 mg in ampoule.
----------------------	---

12.4 Medicines used in heart failure

This subsection will be reviewed at the next meeting of the Expert Committee.

digoxin	Injection: 250 micrograms/ml in 2-ml ampoule. Oral liquid: 50 micrograms/ml. Tablet: 62.5 micrograms; 250 micrograms.
enalapril	Tablet: 2.5 mg.
furosemide	Injection: 10 mg/ml in 2-ml ampoule. Tablet: 40 mg.
hydrochlorothiazide	Tablet (scored): 25 mg.

Complementary List

dopamine	<i>Injection:</i> 40 mg/ml (hydrochloride) in 5-ml vial.
dobutamine (T)	250 mg inj.

12.5 Antithrombotic medicines

acetylsalicylic acid	Tablet: 100 mg.
----------------------	------------------------

Complementary List

streptokinase	<i>Powder for injection:</i> 1.5 million IU in vial.
---------------	--

12.6 Lipid-lowering agents

simvastatin*	Tablet: 5 mg; 10 mg; 20 mg; 40 mg. * For use in high-risk patients.
--------------	---

13. DERMATOLOGICAL MEDICINES (topical)

13.1 Antifungal medicines

benzoic acid + salicylic acid	Ointment or cream: 6% + 3%.
-------------------------------	------------------------------------

©miconazole	Ointment or cream; 2% (nitrate).
sodium thiosulfate	Solution: 15%.

Complementary List

selenium sulfide	Detergent-based suspension: 2%.
------------------	---------------------------------

13.2 Anti-infective medicines

©methylrosanilinium chloride (gentian violet)	Aqueous solution: 0.5%. Tincture: 0.5%.
neomycin sulfate + ©bacitracin	Ointment: 5 mg neomycin sulfate + 250 IU bacitracin zinc/g.
potassium permanganate	Aqueous solution: 1:10 000.
silver sulfadiazine	Cream: 1%, in 500-g container.

13.3 Anti-inflammatory and antipruritic medicines

©betamethasone	Ointment or cream: 0.1% (as valerate).
©calamine lotion	Lotion.
©hydrocortisone	Ointment or cream: 1% (acetate).

13.4 Astringent medicines

aluminium diacetate	Solution: 5%.
---------------------	---------------

13.5 Medicines affecting skin differentiation and proliferation

benzoyl peroxide	Lotion or cream: 5%.
coal tar	Solution: 5%.
dithranol	Ointment: 0.1%-2%.
fluorouracil	Ointment: 5%.
©podophyllum resin	Solution: 10-25%.
salicylic acid	Solution: 5%.
urea	Ointment or cream: 10%.

13.6 Scabicides and pediculicides

©benzyl benzoate	Lotion: 25%.
------------------	--------------

© miconazole	Ointment or cream: 2% (nitrate).
sodium thiosulfate	Solution: 15%.

Complementary List

<i>selenium sulfide</i>	<i>Detergent-based suspension: 2%.</i>
-------------------------	--

13.2 Anti-infective medicines

© methylirosanilinium chloride (gentian violet)	Aqueous solution: 0.5%. Tincture: 0.5%.
neomycin sulfate + © bacitracin	Ointment: 5 mg neomycin sulfate + 250 IU bacitracin zinc/g.
potassium permanganate	Aqueous solution: 1:10 000.
silver sulfadiazine	Cream: 1%, in 500-g container.

13.3 Anti-inflammatory and antipruritic medicines

© betamethasone	Ointment or cream: 0.1% (as valerate).
© calamine lotion	Lotion.
© hydrocortisone	Ointment or cream: 1% (acetate).

13.4 Astringent medicines

aluminium diacetate	Solution: 5%.
---------------------	---------------

13.5 Medicines affecting skin differentiation and proliferation

benzoyl peroxide	Lotion or cream: 5%.
coal tar	Solution: 5%.
dithranol	Ointment: 0.1%-2%.
fluorouracil	Ointment: 5%.
© podophyllum resin	Solution: 10-25%.
salicylic acid	Solution: 5%.
urea	Ointment or cream: 10%.

13.6 Scabicides and pediculicides

© benzyl benzoate	Lotion: 25%.
-------------------	--------------

17. GASTROINTESTINAL MEDICINES

17.1 Antacids and other antiulcer medicines

aluminium hydroxide	Oral liquid: 320 mg/5 ml. Tablet: 500 mg.	@Doctors Care
©ranitidine	Injection: 25 mg/ml in 2-ml ampoule. Oral liquid: 75 mg/5 ml. Tablet: 150 mg (as hydrochloride).	
magnesium hydroxide	Oral liquid: equivalent to 550 mg magnesium oxide/10 ml.	
Lactulose	3.5mg/ 5ml syrup	
Omeprazole	20 mg cap.	

17.2 Antiemetic medicines

metoclopramide	Injection: 5 mg (hydrochloride)/ml in 2-ml ampoule. Tablet: 10 mg (hydrochloride).
promethazine	Injection: 25 mg (hydrochloride)/ml in 2-ml ampoule. Oral liquid: 5 mg (hydrochloride)/5 ml. Tablet: 10 mg; 25 mg (hydrochloride).

17.3 Anti-inflammatory medicines

©sulfasalazine	Retention enema. Suppository: 500 mg. Tablet: 500 mg.
----------------	---

Complementary List

©hydrocortisone	Retention enema. Suppository: 25 mg (acetate). (the ©only applies to hydrocortisone retention enema).
-----------------	--

17.4 Laxatives

©senna	Tablet: 7.5 mg (sennosides) (or traditional dosage forms).
--------	--

17.5 Medicines used in diarrhoea

17.5.1 Oral rehydration

oral rehydration salts*	<p>glucose: 75 mEq sodium: 75 mEq or mmol/l chloride: 65 mEq or mmol/l potassium: 20 mEq or mmol/l citrate: 10 mmol/l osmolarity: 245 mOsm/l glucose: 13.5 g/l sodium chloride: 2.6 g/l potassium chloride: 1.5 g/l trisodium citrate dihydrate+: 2.9 g/l</p> <p>* trisodium citrate dihydrate may be replaced by sodium hydrogen carbonate (sodium bicarbonate) 2.5 g/l. However, as the stability of this latter formulation is very poor under tropical conditions, it is only recommended when manufactured for immediate use.</p> <p>* In cases of cholera a higher concentration of sodium may be required.</p>
-------------------------	---

17.5.2 Medicines for diarrhoea in children

zinc sulfate*	<p>Oral liquid: in 10 mg per unit dosage forms.</p> <p>Tablet: in 10 mg per unit dosage forms.</p> <p>* In acute diarrhoea zinc sulfate should be used as an adjunct to oral rehydration salts.</p>
---------------	---

17.5.3 Antidiarrhoeal (symptomatic) medicines in adults

codeine*	<p>Tablet: 30 mg (phosphate).</p>
----------	-----------------------------------

18. HORMONES, OTHER ENDOCRINE MEDICINES AND CONTRACEPTIVES

18.1 Adrenal hormones and synthetic substitutes

Addison's disease is a rare condition; adrenal hormones are already included in section 3.

18.1 Androgens

Complementary List

testosterone	<p><i>Injection:</i> 200 mg (enantate) in 1-ml ampoule.</p>
--------------	---

18.2 Contraceptives

18.2.1 Oral hormonal contraceptives

©ethinylestradiol + ©levonorgestrel	<p>Tablet: 30 micrograms + 150 micrograms.</p>
--	--

①ethinylestradiol + ②norethisterone	Tablet: 35 micrograms + 1.0 mg.
levonorgestrel	Tablet: 30 micrograms; 750 micrograms (pack of two); 1.5 mg.

18.2.2 Injectable hormonal contraceptives

medroxyprogesterone acetate	Depot injection: 150 mg/ml in 1-ml vial.
medroxyprogesterone acetate + estradiol cypionate	Injection: 25 mg + 5 mg.
norethisterone enantate	Oily solution: 200 mg/ml in 1-ml ampoule.

18.2.3 Intrauterine devices

copper-containing device

18.2.4 Barrier methods

condoms

diaphragms

18.2.5 Implantable contraceptives

levonorgestrel-releasing implant

**@Doctors
Care**

18.3 Estrogens

①ethinylestradiol*	Tablet: 10 micrograms; 50 micrograms. * The public health relevance and/or comparative efficacy and/or safety of this item has been questioned and its continued inclusion on the list will be reviewed at the next meeting of the xpert Committee.
--------------------	--

18.4 Insulins and other antidiabetic agents

glibenclamide	Tablet: 2.5 mg; 5 mg.
insulin injection (soluble)	Injection: 100 IU/ml in 10-ml vial.
intermediate-acting insulin	Injection: 40 IU/ml in 10-ml vial; 100 IU/ml in 10-ml vial (as compound insulin zinc suspension or isophane insulin).
metformin	Tablet: 500 mg (hydrochloride).

18.5 Ovulation inducers

Complementary List

clomifene

Tablet: 50 mg (citrate).

18.6 Progestogens

norethisterone*	Tablet: 5 mg. * The public health relevance and/or comparative efficacy and/or safety of this item has been questioned and its continued inclusion on the list will be reviewed at the next meeting of the expert Committee.
-----------------	---

Complementary List

medroxyprogesterone acetate*

Tablet: 5 mg.

* The public health relevance and/or comparative efficacy and/or safety of this item has been questioned and its continued inclusion on the list will be reviewed at the next meeting of the expert Committee.

18.7 Thyroid hormones and antithyroid medicines

levothyroxine	Tablet: 50 micrograms; 100 micrograms (sodium salt).
potassium iodide	Tablet: 60 mg.
© propylthiouracil	Tablet: 50 mg.

19. IMMUNOLOGICALS**19.1 Diagnostic agents**

tuberculin, purified protein derivative (PPD)	Injection.
---	------------

19.2 Sera and immunoglobulins

All plasma fractions should comply with the WHO Requirements for the Collection, Processing and Quality Control of Blood, Blood Components and Plasma Derivatives.

anti-D immunoglobulin (human)	Injection: 250 micrograms in single-dose vial.
antitetanus immunoglobulin (human)	Injection: 500 IU in vial.
antivenom immunoglobulin*	Injection. * Exact type to be defined locally.
diphtheria antitoxin	Injection: 10 000 IU; 20 000 IU in vial.

**@Doctors
Care**

© rabies immunoglobulin	Injection: 150 IU/ml in vial.
Anti hepatitis B immunoglobulin	injectin.
19.3 Vaccines	
Vaccine for Universal Immunization.	
BCG (dried) vaccine	inj
diphtheria-pertussis tetanus vaccine	inj.
diphtheria-tetanus vaccine	inj.
measles-mumps-rubella vaccine	
measles vaccine	inj.
poliomyelitis vaccine	oral solution, live attenuated
tetanus vaccine	inj.
Haemophilus influenzae type B vaccine	inj.
Rota virus vaccine	inj.
Typhoid Vaccines	inj.
Vaccines for Specific Use.	
Hepatitis A vaccine	inj
hepatitis B vaccine	inj.
meningococcal vaccine	inj.
rabies vaccine	inj.
rubella vaccine	inj
Cholera Vaccines	inj
Flu vaccine (influenza)	inj
Pneumococcal vaccine	inj

@Doctors
Care

20. MUSCLE RELAXANTS (PERIPHERALLY-ACTING) AND CHOLINESTERASE INHIBITORS

©alcuronium	<i>Injection:</i> 5 mg (chloride)/ml in 2-ml ampoule.
neostigmine	<i>Injection:</i> 500 micrograms in 1-ml ampoule; 2.5 mg (metilsulfate) in 1-ml ampoule. <i>Tablet:</i> 15 mg (bromide).
suxamethonium	<i>Injection:</i> 50 mg (chloride)/ml in 2-ml ampoule. <i>Powder for injection</i> (chloride) in vial.

Complementary List

pyridostigmine	<i>Injection:</i> 1 mg in 1-ml ampoule. <i>Tablet:</i> 60 mg (bromide).
©vecuronium	<i>Powder for injection:</i> 10 mg (bromide) in vial.

21. OPHTHALMOLOGICAL PREPARATIONS

This section will be reviewed at the next meeting of the Expert Committee.

21.1 Anti-infective agents

aciclovir	Ointment: 3% W/W.
©gentamicin*	Solution (eye drops): 0.3% (sulfate). * Final selection depends on indication for use.
©tetracycline	Eye ointment: 1% (hydrochloride).

21.2 Anti-inflammatory agents

©prednisolone	Solution (eye drops): 0.5% (sodium phosphate).
---------------	---

21.3 Local anaesthetics

©tetracaine	Solution (eye drops): 0.5% (hydrochloride).
-------------	--

21.4 Miotics and antiglaucoma medicines

acetazolamide	Tablet: 250 mg.
©pilocarpine	Solution (eye drops): 2%; 4% (hydrochloride or nitrate).
©timolol	Solution (eye drops): 0.25%; 0.5% (as maleate).

21.5 Mydriatics

atropine	Solution (eye drops): 0.1%; 0.5%, 1% (sulfate).
----------	--

Complementary List

<i>epinephrine (adrenaline)</i>	<i>Solution (eye drops): 2% (as hydrochloride).</i>
22. OXYTOCICS AND ANTIOXYTOCICS	
22.1 Oxytocics	
ergometrine	<i>Injection: 200 micrograms (hydrogen maleate) in 1-ml ampoule.</i>
oxytocin	<i>Injection: 10 IU in 1-ml ampoule.</i>
22.2 Antioxytocics (tocolytics)	
nifedipine	<i>Immediate release capsule: 10 mg.</i>

23. PERITONEAL DIALYSIS SOLUTION	
<i>Complementary List</i>	
<i>intraperitoneal dialysis solution (of appropriate composition)</i>	<i>PARENTERAL solution.</i>
24. PSYCHOTHERAPEUTIC MEDICINES	
24.1 Medicines used in psychotic disorders	
chlorpromazine	<i>Injection: 25 mg (hydrochloride)/ml in 2-ml ampoule. Oral liquid: 25 mg (hydrochloride)/5 ml. Tablet: 100 mg (hydrochloride).</i>
fluphenazine	<i>Injection: 25 mg (decanoate or enantate) in 1-ml ampoule.</i>
haloperidol	<i>Injection: 5 mg in 1-ml ampoule. Tablet: 2 mg; 5 mg.</i>
24.2 Medicines used in mood disorders	
24.2.1 Medicines used in depressive disorders	
amitriptyline	<i>Tablet: 25 mg (hydrochloride).</i>
fluoxetine	<i>Capsule or tablet: 20 mg (present as hydrochloride).</i>
24.2.2 Medicines used in bipolar disorders	
carbamazepine	<i>Tablet (scored): 100 mg; 200 mg.</i>
lithium carbonate	<i>Capsule or tablet: 300 mg.</i>

valproic acid	Tablet (enteric-coated): 200 mg; 500 mg (sodium valproate).
Risperidone	1 mg, 2 mg, 3 mg tab.
24.3 Medicines used in generalized anxiety and sleep disorders	
©diazepam	Tablet (scored): 2 mg; 5 mg.
24.4 Medicines used for obsessive compulsive disorders and panic attacks	
clomipramine	Capsule: 10 mg; 25 mg (hydrochloride).
24.5 Medicines used in substance dependence programmes	
<i>Complementary List</i>	
©methadone*	<p><i>Concentrate for oral liquid:</i> 5 mg/ml; 10 mg/ml (hydrochloride).</p> <p><i>Oral liquid:</i> 5 mg/5 ml; 10 mg/5 ml.</p> <p>* The square box is added to include buprenorphine. The medicines should only be used within an established support programme.</p>
25. MEDICINES ACTING ON THE RESPIRATORY TRACT	
25.1 Antiasthmatic and medicines for chronic obstructive pulmonary disease	
©beclometasone	Inhalation (aerosol): 50 micrograms per dose (dipropionate); 250 micrograms (dipropionate) per dose.
epinephrine (adrenaline)	Injection: 1 mg (as hydrochloride or hydrogen tartrate) in 1-ml ampoule.
ipratropium bromide	Inhalation (aerosol): 20 micrograms/metered dose.
©salbutamol	<p>Inhalation (aerosol): 100 micrograms (as sulfate) per dose.</p> <p>Injection: 50 micrograms (as sulfate)/ml in 5-ml ampoule.</p> <p>Oral liquid: 2 mg/5 mL</p> <p>Respirator solution for use in nebulizers: 5 mg (as sulfate)/ml.</p> <p>Tablet: 2 mg; 4 mg (as sulfate).</p>

25.2 Other medicines acting on the respiratory tract

**@Doctors
Care**